AMENDMENTS TO THE CLAIMS

- 20. (Currently amended) An isolated nucleic acid molecule <u>consisting of comprising</u> a polynucleotide sequence selected from the group consisting of:
 - (a) an isolated polynucleotide encoding a polypeptide corresponding to amino acids 1 to 325 of SEQ ID NO:2 including the start codon;
 - (b) an isolated polynucleotide encoding a polypeptide corresponding to amino acids 2 to 325 of SEQ ID NO:2 minus the start codon;
 - (c) an isolated polynucleotide encoding a polypeptide corresponding to amino acids 34 to 134 of SEQ ID NO:2;
 - (d) an isolated polynucleotide encoding the K+betaM6 polypeptide as encoded by the cDNA clone contained in ATCC Deposit No: PTA-3161;
 - (e) an isolated polynucleotide encoding at least 50 contiguous amino acids of SEQ ID NO:2;
 - (f) an isolated polynucleotide which represents the complimentary sequence (antisense) of (a), (b), (c), (d), (e) or \underline{a} fragment thereof; and
 - (g) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(f), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.
- 21. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (a).
- 22. (Original) The isolated nucleic acid molecule of claim 21, wherein said polynucleotide comprises nucleotides 121 to 1095 of SEQ ID NO:1.
- 23. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (b).

- 24. (Original) The isolated nucleic acid molecule of claim 23, wherein said polynucleotide comprises nucleotides 124 to 1095 of SEQ ID NO:1.
- 25. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (c).
- 26. (Original) The isolated nucleic acid molecule of claim 25, wherein said polynucleotide comprises nucleotides 220 to 522 of SEQ ID NO:1.
- 27. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (d).
- 28. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (e).
- 29. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (f).
- 30. (Original) The isolated nucleic acid molecule of claim 20, wherein said polynucleotide is (g).
- 31. (Original) A recombinant vector comprising the isolated nucleic acid molecule of claim 20.
 - 32. (Original) A recombinant host cell comprising the vector sequences of claim 31.
 - 33. (Original) A method of making an isolated polypeptide comprising:
 - (a) culturing the recombinant host cell of claim 32 under conditions such that said polypeptide is expressed; and
 - (b) recovering said polypeptide.

- 34. (Original) A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:
 - (a) determining the presence or absence of a mutation in the polynucleotide of claim 20; and
 - (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.
- 35. (Original) The isolated polynucleotide of claim 20 wherein said nucleic acid sequence further comprises a heterologous nucleic acid sequence.
- 36. (Original) The isolated polynucleotide of claim 35 wherein said heterologous nucleic acid sequence encodes a heterologous polypeptide.
- 37. (Original) The isolated polynucleotide of claim 36 wherein said heterologous polypeptide is the Fc domain of immunoglobulin.
- 38. (Original) An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 70.0% identical to a sequence provided in claim 20, wherein percent identity is calculated using a CLUSTALW global sequence alignment.
- 39. (Original) The isolated polynucleotide of claim 20 wherein said nucleic acid sequence further comprises a heterologous nucleic acid sequence.
- 40. (Original) The isolated polynucleotide of claim 35 wherein said heterologous nucleic acid sequence encodes a heterologous polypeptide.
- 41. (Original) The isolated polynucleotide of claim 36 wherein said heterologous polypeptide is the Fc domain of immunoglobulin.